

# **PRK Post-Operative**

### **MEDICATED DROPS:**

VIGAMOX (moxifloxacin): This is an antibiotic in order to prevent infections.

One drop to the operated eye **<u>3timesperday</u> x <u>1week</u>** 

<u>PRED FORTE (prednisolone acetate)</u>: This is an anti-inflammatory medication to help the eye heal. One drop to the operated eye <u>3timesperday</u> x 2 weeks, then 2 times per day x 1 week, 1 times per day x 1 week making for <u>4 weeks total</u>

**<u>BROMDAY</u>** (bromfenac): This is an anti-inflammatory medication to help the eye heal and for pain relief. One drop to the operated eye <u>**2timesperday</u>** for 2-3 days only</u>

**DOXYCYCLINE:** This is an antibiotic with anti-inflammatory action preventing corneal scarring. **Take 100mg daily** for 14 days

VITAMIN C: This is a vitamin that helps the cornea heal faster. Take 1000mg daily for 1 month

**VOLTAREN:** This is an anti-inflammatory pill for pain control. **Take 100mg 1time per day with meals ONLY if there is pain** 

	PRED FORTE BORNES BORNE	VIGAMOX	BROMDAY
Week 1	3x/day	3x/day	2x/day (only for 2 days)
Week 2	3x/day		
Week 3	2x/day		
Week 4	1x/day		

### **Medication Instructions**

- Wait 2-5 minutes between drops so that the medications do not wash each other out.
- It is not necessary to interrupt your sleep (nap or night) for the medicated drop schedule.
- Follow physician directives on frequency and timing of drops if other than listed above.

Artificial Tears (not pictured) twist top/single use:

- Use only preservative-free eye drops during the first week following surgery. Subsequently, you may use preserved drops that are available in larger, multi-use bottles.
- <u>Administering Drops</u>: Do not lift top eyelid. You may pull down gently on lower lid if necessary to put drops in.
- <u>Usage</u>: May start using day of or day after surgery. Use at least once an hour (while awake), but may use more often as desired.

#### **Additional Instructions:**

- Wear provided clear eye goggles at night for 5 days.
- Wear provided dark glasses or high-quality sunglasses in bright sunlight,
  - overly bright rooms, or on windy days for first week.
- Do not rub, bump, poke, scratch, or touch your eyes.

<u>Following Surgery</u>: After your surgery today, expect some discomfort such as a foreign body sensation, burning, stinging, red spots, and/or watering. These sensations are normal and usually resolve within 2-4 days (each patient varies).

- Avoid rubbing, bumping, or scratching the eye; instead use lubricant drops for irritation. <u>Note</u>: The desire to rub your eyes will be a natural instinct; however, you need to wait a few weeks for even the most gentle and light rubbing. For now, **dab and pat only**.
- Rest your eye(s) for 3-4 hours following the procedure and avoid reading, using computers, and watching television on the day of surgery.
- Avoid alcohol.
- No swimming or use of hot tubs.
- Wear sunglasses when outdoors.

#### General Instructions:

- **Headache** Use over-the-counter medications like Advil or Ibuprofen. Do not take Tylenol in conjunction with Vicodin.
- Use a bath, not shower for the first 24 hours in order to avoid soap, shampoo, or water getting in your eyes.
- Showering is okay on the day after surgery. Avoid eye contact with direct flow from the shower head.
- **Driving** is okay once the surgeon clears you. Generally, this is after the contact lens removal appointment.
- Avoid areas or situations with a lot of dirt, dust, debris, pet hair, fuzz, or lint. Note: If you encounter these things, do not rub the eye. Flush with lubricant.
- **Fluctuation** It is normal for your vision to fluctuate slightly until the cornea heals. Increased use of lubricating drops could help reduce fluctuation. This can occur for up to 1-2 months post **PRK**.
- You may begin wearing eye make-up (i.e., eye shadow & mascara) 2 weeks after **PRK**.
- Complete all post-operative care as instructed.

24 Hours after Surgery:

- Reading, watching television, and using a computer are acceptable as long as adequate eye lubrication is maintained.
- Flying in an airplane is all right, but keep the eye generously lubricated (every 30 minutes) since airplanes have very dry air.
- Driving may be resumed if adequate vision is confirmed at the 5-6 day contact lens removal appointment.
- When showering, keep your eyes closed while washing face and hair. Do not rub your eyes when toweling off.

One Week after Surgery and Beyond:

• You may resume normal activities with caution, paying attention to any potential for eye injury. For extreme sports, extreme environments, or unusual activities, please consult with your eye doctor.

Surgery Counselor:

By signing above, I am indicating that I have reviewed pre- and post-operative instructions with the patient and I have given the patient an opportunity to ask questions prior to surgery.

# After-Hours Emergency Number: 917-460-7065